003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

MPG Polyclonal Antibody

Cat #: G-AB-00834

Host: Rabbit **Reactivity:** Human

Isotype: IgG **Clonality:** Polyclonal

Overview:
Maintenance of DNA sequences is necessary for vertebrates and other life. DNA is under constant stress by a plethora of DNA-damaging agents present in both the environment and within cells. The potentially deleterious effects of DNA lesions in cells are elegantly resolved by sophisticated DNA repair systems, including base excision repair (BER), nucleotide excision repair (NER) and DNA repair methyltransferase (MTase). Methylated bases, such as 3-methyladenine (3MeA) and 7-methylguanine (7MeG) can be formed by agents in the environment and by endogenous cellular processes. Consequently, in the absence of exposure to environmental agents, DNA methylation damage can be incurred on the genomic DNA of normal mammalian cells. DNA N-glycosylases are base excision-repair proteins that locate and cleave damaged bases from DNA as the first step in restoring the sequence.

Gene ID:

Accession #: BC014991

Immunogen: Recombinant protein of human MPG

Conjugation: Unconjugated

Swissprot: P29372

Calculated Molecular Weight: 32 kDa

Observed Molecular Weight:

Concentration: 0.5 mg/mL

Buffer: PBS with 0.05% sodium azide and 50% glycerol, PH7.4

Purification Method: Affinity purification

Application: WB,IHC,ELISA

Dilution: WB 1:1000-1:5000, IHC 1:50-1:200

Storage: Store at -20°C. Avoid freeze / thaw cycles.