003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

# PTPRM Polyclonal Antibody

**Cat #:** G-AB-01478

**Host:** Rabbit **Reactivity:** Human, Mouse

**Isotype:** IgG **Clonality:** Polyclonal

**Overview:**
The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP possesses an extracellular region, a single transmembrane region, and two tandem catalytic domains, and thus represents a receptor-type PTP. The extracellular region contains a meprin-A5 antigen-PTP mu (MAM) domain, an Ig-like domain and four fibronectin type III-like repeats. This PTP has been shown to mediate cell-cell aggregation through the interaction with another molecule of this PTP on an adjacent cell. This PTP can interact with scaffolding protein RACK1/GNB2L1, whicHuman, Mouseay be necessary for the downstream signaling in response to cell-cell adhesion. Alternative splicing results in multiple transcripts encoding distinct isoforms.

**Gene ID:**

**Accession #:** NP_002836

**Immunogen:** Synthetic peptide of human PTPRM

**Conjugation:** Unconjugated

**Swissprot:** P28827

**Calculated Molecular Weight:**

**Observed Molecular Weight:**

**Concentration:** 0.4 mg/mL

**Buffer:** PBS with 0.05% sodium azide and 50% glycerol, PH7.4

**Purification Method:** Affinity purification

**Application:** IHC,ELISA

**Dilution:** IHC 1:50-1:200

**Storage:** Store at -20°C. Avoid freeze / thaw cycles.