003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

MT-ND1 Polyclonal Antibody

Cat #: G-AB-01779

Host: Rabbit **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:** Polyclonal

Overview:
NADH:ubiquinone oxidoreductase (complex I) is an extremely complicated multiprotein complex located in the inner mitochondrial membrane. Human complex I is important for energy metabolism because its main function is to transport electrons from NADH to ubiquinone, which is accompanied by translocation of protons from the mitochondrial matrix to the intermembrane space. Human complex I appears to consist of 41 subunits. A small number of complex I subunits are the products of mitochondrial genes (subunits 1-7), while the remainder are nuclear encoded and imported from the cytoplasm. NADH dehydrogenase subunit 1 (ND1) binds rotenone and rotenone analogs and might be involved in electron transfer to ubiquinone. Mutations in the ND1 gene may be implicated in several disorders, including Leber hereditary optic neuropathy, Alzheimer disease, and Parkinson disease.

Gene ID:

Accession #: YP_003024026

Immunogen: Synthetic peptide of human MT-ND1

Conjugation: Unconjugated

Swissprot: P03886

Calculated Molecular Weight: 36 kDa

Observed Molecular Weight:

Concentration: 1 mg/mL

Buffer: PBS with 0.05% sodium azide and 50% glycerol, PH7.4

Purification Method: Affinity purification

Application: WB,IHC,ELISA

Dilution: WB 1:200-1:1000, IHC 1:50-1:200

Storage: Store at -20°C. Avoid freeze / thaw cycles.