003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

ERG Polyclonal Antibody

Cat #: G-AB-02028

Host: Rabbit **Reactivity:** Human, Mouse

Isotype: IgG **Clonality:** Polyclonal

Overview:
This gene encodes a member of the erythroblast transformation-specific (ETS) family of transcriptions factors. All members of this family are key regulators of embryonic development, cell proliferation, differentiation, angiogenesis, inflammation, and apoptosis. The protein encoded by this gene is mainly expressed in the nucleus. It contains an ETS DNA-binding domain and a PNT (pointed) domain which is implicated in the self-association of chimeric oncoproteins. This protein is required for platelet adhesion to the subendothelium, inducing vascular cell remodeling. It also regulates hematopoesis, and the differentiation and maturation of megakaryocytic cells. This gene is involved in chromosomal translocations, resulting in different fusion gene products, such as TMPSSR2-ERG and NDRG1-ERG in prostate cancer, EWS-ERG in Ewing's sarcoma and FUS-ERG in acute myeloid leukemia. Multiple alternatively spliced transcript variants encoding different isoforms have been identified.

Gene ID:

Accession #: BC040168

Immunogen: Recombinant protein of human ERG

Conjugation: Unconjugated

Swissprot: P11308

Calculated Molecular Weight: 54 kDa

Observed Molecular Weight:

Concentration: 0.4 mg/mL

Buffer: PBS with 0.05% sodium azide and 50% glycerol, PH7.4

Purification Method: Affinity purification

Application: WB,IHC,ELISA

Dilution: WB 1:500-1:2000, IHC 1:50-1:200

Storage: Store at -20°C. Avoid freeze / thaw cycles.