003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

CD10 Polyclonal Antibody

Cat #: G-AB-02893

Host: Rabbit **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:** Polyclonal

Overview:
This gene encodes a common acute lymphocytic leukemia antigen that is an important cell surface marker in the diagnosis of human acute lymphocytic leukemia (ALL). This protein is present on leukemic cells of pre-B phenotype, which represent 85% of cases of ALL. This protein is not restricted to leukemic cells, however, and is found on a variety of normal tissues. It is a glycoprotein that is particularly abundant in kidney, where it is present on the brush border of proximal tubules and on glomerular epithelium. The protein is a neutral endopeptidase that cleaves peptides at the amino side of hydrophobic residues and inactivates several peptide hormones including glucagon, enkephalins, substance P, neurotensin, oxytocin, and bradykinin. This gene, which encodes a 100-kD type II transmembrane glycoprotein, exists in a single copy of greater than 45 kb. The 5' untranslated region of this gene is alternatively spliced, resulting in four separate mRNA transcripts. The coding region is not affected by alternative splicing.

Gene ID:

Accession #: NP_000893

Immunogen: Synthetic peptide of human MME

Conjugation: Unconjugated

Swissprot: P08473

Calculated Molecular Weight:

Observed Molecular Weight:

Concentration: 0.5 mg/mL

Buffer: PBS with 0.05% sodium azide and 50% glycerol, PH7.4

Purification Method: Affinity purification

Application: IHC,ELISA

Dilution: IHC 1:50-1:200

Storage: Store at -20°C. Avoid freeze / thaw cycles.