003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

LCN2 Polyclonal Antibody

Cat #: G-AB-02899

Host: Rabbit **Reactivity:** Human

Isotype: IgG **Clonality:** Polyclonal

Overview:
Iron-trafficking protein involved in multiple processes such as apoptosis, innate immunity and renal development. Binds iron through association with 2,5-dihydroxybenzoic acid (2,5-DHBA), a siderophore that shares structural similarities with bacterial enterobactin, and delivers or removes iron from the cell, depending on the context. Iron-bound form (holo-24p3) is internalized following binding to the SLC22A17 (24p3R) receptor, leading to release of iron and subsequent increase of intracellular iron concentration. In contrast, association of the iron-free form (apo-24p3) with the SLC22A17 (24p3R) receptor is followed by association with an intracellular siderophore, iron chelation and iron transfer to the extracellular medium, thereby reducing intracellular iron concentration. Involved in apoptosis due to interleukin-3 (IL3) deprivation: iron-loaded form increases intracellular iron concentration without promoting apoptosis, while iron-free form decreases intracellular iron levels, inducing expression of the proapoptotic protein BCL2L11/BIM, resulting in apoptosis. Involved in innate immunity, possibly by sequestrating iron, leading

Gene ID:

Accession #: NP_005555

Immunogen: Synthetic peptide of human LCN2

Conjugation: Unconjugated

Swissprot: P80188

Calculated Molecular Weight: 23 kDa

Observed Molecular Weight:

Concentration: 0.6 mg/mL

Buffer: PBS with 0.05% sodium azide and 50% glycerol, PH7.4

Purification Method: Affinity purification

Application: WB,IHC,ELISA

Dilution: WB 1:1000-1:5000, IHC 1:50-1:200

Storage: Store at -20°C. Avoid freeze / thaw cycles.