003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

SMC1A Polyclonal Antibody

Cat #: G-AB-03949

Host: Rabbit **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:** Polyclonal

Overview:
Proper cohesion of sister chromatids is a prerequisite for the correct segregation of chromosomes during cell division. The cohesin multiprotein complex is required for sister chromatid cohesion. This complex is composed partly of two structural maintenance of chromosomes (SMC) proteins, SMC3 and either SMC1B or the protein encoded by this gene. Most of the cohesin complexes dissociate from the chromosomes before mitosis, although those complexes at the kinetochore remain. Therefore, the encoded protein is thought to be an important part of functional kinetochores. In addition, this protein interacts with BRCA1 and is phosphorylated by ATM, indicating a potential role for this protein in DNA repair. This gene, which belongs to the SMC gene family, is located in an area of the X-chromosome that escapes X inactivation. Mutations in this gene result in Cornelia de Lange syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms.

Gene ID:

Accession #: BC112127

Immunogen: Fusion protein of human SMC1A

Conjugation: Unconjugated

Swissprot: Q14683

Calculated Molecular Weight:

Observed Molecular Weight:

Concentration: 0.7 mg/mL

Buffer: PBS with 0.05% NaN3 and 40% Glycerol, pH7.4

Purification Method: Antigen affinity purification

Application: IHC,ELISA

Dilution: IHC 1:30-1:150, ELISA 1:5000-1:10000

Storage: Store at -20°C. Avoid freeze / thaw cycles.