003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

CCL21 Polyclonal Antibody

Cat #: G-AB-04725

Host: Rabbit **Reactivity:** Human

Isotype: IgG **Clonality:** Polyclonal

Overview:
This gene is one of several CC cytokine genes clustered on the p-arm of chromosome 9. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. Similar to other chemokines the protein encoded by this gene inhibits hemopoiesis and stimulates chemotaxis. This protein is chemotactic in vitro for thymocytes and activated T cells, but not for B cells, macrophages, or neutrophils. The cytokine encoded by this gene may also play a role in mediating homing of lymphocytes to secondary lymphoid organs. It is a high affinity functional ligand for chemokine receptor 7 that is expressed on T and B lymphocytes and a known receptor for another member of the cytokine family (small inducible cytokine A19).

Gene ID:

Accession #: NP002980

Immunogen: Synthetic peptide of human CCL21

Conjugation: Unconjugated

Swissprot: O00585

Calculated Molecular Weight: 15 kDa

Observed Molecular Weight: Refer to figures

Concentration: 0.9 mg/mL

Buffer: PBS with 0.05% NaN3 and 40% Glycerol, pH7.4

Purification Method: Antigen affinity purification

Application: WB,ELISA

Dilution: WB 1:500-1:2000, ELISA 1:5000-1:10000

Storage: Store at -20°C. Avoid freeze / thaw cycles.