003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

C14orf169 Polyclonal Antibody

Cat #: G-AB-04832

Host: Rabbit **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:** Polyclonal

Overview:
Oxygenase that can act as both a histone lysine demethylase and a ribosomal histidine hydroxylase. Specifically demethylates 'Lys-4' (H3K4me) and 'Lys-36' (H3K36me) of histone H3, thereby playing a central role in histone code. Preferentially demethylates trimethylated H3 'Lys-4' (H3K4me3) and monomethylated H3 'Lys-4' (H3K4me1) residues, while it has weaker activity for dimethylated H3 'Lys-36' (H3K36me2). Also catalyzes the hydroxylation of 60S ribosomal protein L8 on 'His-216'. Acts as a regulator of osteoblast differentiation via its interaction with SP7/OSX by demethylating H3K4me and H3K36me, thereby inhibiting SP7/OSX-mediated promoter activation (By similarity). May also play a role in ribosome biogenesis and in the replication or remodeling of certain heterochromatic region. Participates in MYC-induced transcriptional activation.

Gene ID:

Accession #: NP078920

Immunogen: Synthetic peptide of human C14orf169

Conjugation: Unconjugated

Swissprot: Q9H6W3

Calculated Molecular Weight:

Observed Molecular Weight:

Concentration: 0.8 mg/mL

Buffer: PBS with 0.05% NaN3 and 40% Glycerol, pH7.4

Purification Method: Antigen affinity purification

Application: IHC,ELISA

Dilution: IHC 1:30-1:150, ELISA 1:5000-1:10000

Storage: Store at -20°C. Avoid freeze / thaw cycles.