003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

HMGB1 Polyclonal Antibody

Cat #: G-AB-05211

Host: Rabbit **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:** Polyclonal

Overview:
High mobility group (HMG) proteins 1 and 2 are ubiquitous non-histone components of chromatin. Evidence suggests that the binding of HMG proteins to DNA induces alterations in the DNA architecture including DNA bending and unwinding of the helix. HMG proteins synergize with Oct-2, members of the NFB family, ATF-2 and c-Jun to activate transcription. Other studies indicate that phosphorylation of HMG protein is required to stimulate the transcriptional activity of the protein. Human HMG-1 and HMG-2 both contain two DNA-binding domains, termed HMG boxes. HMG proteins bind single-stranded DNA but induce conformational changes in double-stranded DNA alone.

Gene ID:

Accession #:

Immunogen: KLH conjugated Synthetic peptide corresponding to Mouse HMGB1

Conjugation: Unconjugated

Swissprot: P09429,P63158,P63159

Calculated Molecular Weight: 25 kDa

Observed Molecular Weight: 25 kDa

Concentration: 1.45 mg/mL

Buffer: PBS with 0.02% sodium azide, 1% BSA and 50% glycerol, pH7.4

Purification Method: Affinity purification

Application: WB,IHC,IF

Dilution: WB 1:500-1:2000, IHC 1:200-1:800, IF 1:200-1:800

Storage: Store at -20°C. Avoid freeze / thaw cycles.