003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

STAT1 Monoclonal Antibody

Cat #: G-AB-05675

Host: Mouse **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:** Monoclonal

Overview:
Signal transducer and activator of transcription that mediates signaling by interferons (IFNs). Following type I IFN (IFN-alpha and IFN-beta) binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize, associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of interferon stimulated genes, which drive the cell in an antiviral state. In response to type II IFN (IFN-gamma), STAT1 is tyrosine- and serine-phosphorylated. It then forms a homodimer termed IFN-gamma-activated factor (GAF), migrates into the nucleus and binds to the IFN gamma activated sequence (GAS) to drive the expression of the target genes, inducing a cellular antiviral state.

Gene ID:

Accession #:

Immunogen: Synthetic Peptide of STAT1

Conjugation: Unconjugated

Swissprot: P42224

Calculated Molecular Weight:

Observed Molecular Weight:

Concentration: 1 mg/mL

Buffer: PBS with 0.02% sodium azide, 0.5% BSA and 50% glycerol, pH7.4

Purification Method: Protein A purification

Application: IHC-p

Dilution: IHC 1:100-200

Storage: Store at -20°C. Avoid freeze / thaw cycles.