003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

KCTD16 Polyclonal Antibody

Cat #: G-AB-06202

Host: Rabbit **Reactivity:** Human, Mouse

Isotype: IgG **Clonality:** Polyclonal

Overview:
The BTB (Broad-Complex, Tramtrack and Bric a brac) domain, also known as the POZ (Poxvirus and Zinc finger) domain, is an N-terminal homodimerization domain that contains multiple copies of kelch repeats and/or C2H2-type zinc fingers. Proteins that contain BTB domains are thought to be involved in transcriptional regulation via control of chromatin structure and function. KCTD16 (potassium channel tetramerisation domain containing 16), also known as BTB/POZ domain-containing protein KCTD16, is a 428 amino acid protein that contains one BTB (POZ) domain. An auxiliary subunit of GABAB R1 and GABAB R2, KCTD16 increases agonist potency and alters the G-protein signaling of the receptors by accelerating onset and promoting desensitization.

Gene ID:

Accession #: BC113435

Immunogen: Fusion protein of human KCTD16

Conjugation: Unconjugated

Swissprot: Q68DU8

Calculated Molecular Weight: 49 kDa

Observed Molecular Weight: Refer to figures

Concentration: 0.8 mg/mL

Buffer: PBS with 0.05% NaN3 and 40% Glycerol, pH7.4

Purification Method: Antigen affinity purification

Application: WB,IHC,ELISA

Dilution: WB 1:500-1:2000, IHC 1:25-1:100, ELISA 1:5000-1:10000

Storage: Store at -20°C. Avoid freeze / thaw cycles.