003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

METAP1D Polyclonal Antibody

Cat #: G-AB-06203

Host: Rabbit **Reactivity:** Human, Mouse

Isotype: IgG **Clonality:** Polyclonal

Overview:
The N-terminal methionine excision pathway is an essential process in which the N-terminal methionine is removed from many proteins, thus facilitating subsequent protein modification. In mitochondria, enzymes that catalyze this reaction are celled methionine aminopeptidases (MetAps, or MAPs; EC 3.4.11.18) (Serero et al., 2003 [PubMed 14532271])

Gene ID:

Accession #: BC113644

Immunogen: Fusion protein of human METAP1D

Conjugation: Unconjugated

Swissprot: Q6UB28

Calculated Molecular Weight: 37 kDa

Observed Molecular Weight: Refer to figures

Concentration: 0.9 mg/mL

Buffer: PBS with 0.05% NaN3 and 40% Glycerol, pH7.4

Purification Method: Antigen affinity purification

Application: WB,IHC,ELISA

Dilution: WB 1:500-1:2000, IHC 1:40-1:200, ELISA 1:5000-1:10000

Storage: Store at -20°C. Avoid freeze / thaw cycles.