003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

PSMD12 Polyclonal Antibody

Cat #: G-AB-06489

Host: Rabbit **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:** Polyclonal

Overview:
The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator. A pseudogene has been identified on chromosome 3. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Gene ID:

Accession #: BC019062

Immunogen: Fusion protein of human PSMD12

Conjugation: Unconjugated

Swissprot: O00232

Calculated Molecular Weight: 53 kDa

Observed Molecular Weight: Refer to figures

Concentration: 0.9 mg/mL

Buffer: PBS with 0.05% NaN3 and 40% Glycerol, pH7.4

Purification Method: Antigen affinity purification

Application: WB,IHC,ELISA

Dilution: WB 1:1000-1:5000, IHC 1:50-1:300, ELISA 1:5000-1:10000

Storage: Store at -20°C. Avoid freeze / thaw cycles.