003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

PNRC2 Polyclonal Antibody

Cat #: G-AB-06540

Host: Rabbit **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:** Polyclonal

Overview:
Involved in nonsense-mediated mRNA decay (NMD) by acting as a bridge between the mRNA decapping complex and the NMD machinery (PubMed:19150429). May act by targeting the NMD machinery to the P-body and recruiting the decapping machinery to aberrant mRNAs (PubMed:19150429). Required for UPF1/RENT1 localization to the P-body (PubMed:19150429). Plays a role in glucocorticoid receptor-mediated mRNA degradation by interacting with the glucocorticoid receptor NR3C1 in a ligand-dependent manner when it is bound to the 5' UTR of target mRNAs and recruiting the RNA helicase UPF1 and the mRNA-decapping enzyme DCP1A, leading to RNA decay (PubMed:25775514). Also acts as a nuclear receptor coactivator (PubMed:11574675). May play a role in controlling the energy balance between energy storage and energy expenditure (By similarity).

Gene ID:

Accession #: BC001959

Immunogen: Fusion protein of human PNRC2

Conjugation: Unconjugated

Swissprot: Q9NPJ4

Calculated Molecular Weight:

Observed Molecular Weight:

Concentration: 0.84 mg/mL

Buffer: PBS with 0.05% NaN3 and 40% Glycerol, pH7.4

Purification Method: Antigen affinity purification

Application: IHC,ELISA

Dilution: IHC 1:50-1:300, ELISA 1:5000-1:10000

Storage: Store at -20°C. Avoid freeze / thaw cycles.