003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

CHEK2 Polyclonal Antibody

Cat #: G-AB-06931

Host: Rabbit **Reactivity:** Human

Isotype: IgG **Clonality:** Polyclonal

Overview:
In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Also, mutations in this gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors. This nuclear protein is a member of the CDS1 subfamily of serine/threonine protein

Gene ID:

Accession #: NP001005735

Immunogen: Synthetic peptide of human CHEK2

Conjugation: Unconjugated

Swissprot: O96017

Calculated Molecular Weight: 61 kDa

Observed Molecular Weight: Refer to figures

Concentration: 0.66 mg/mL

Buffer: PBS with 0.05% NaN3 and 40% Glycerol, pH7.4

Purification Method: Antigen affinity purification

Application: WB,ELISA

Dilution: WB 1:500-1:2000, ELISA 1:5000-1:10000

Storage: Store at -20°C. Avoid freeze / thaw cycles.