003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

SLC27A2 Polyclonal Antibody

Cat #: G-AB-07440

Host: Rabbit **Reactivity:** Human, Mouse

Isotype: IgG **Clonality:**

Overview:
The protein encoded by this gene is an isozyme of long-chain fatty-acid-coenzyme A ligase family. Although differing in substrate specificity, subcellular localization, and tissue distribution, all isozymes of this family convert free long-chain fatty acids into fatty acyl-CoA esters, and thereby play a key role in lipid biosynthesis and fatty acid degradation. This isozyme activates long-chain, branched-chain and very-long-chain fatty acids containing 22 or more carbons to their CoA derivatives. It is expressed primarily in liver and kidney, and is present in both endoplasmic reticulum and peroxisomes, but not in mitochondria. Its decreased peroxisomal enzyme activity is in part responsible for the biochemical pathology in X-linked adrenoleukodystrophy. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Gene ID: 11001

Accession #:

Immunogen: Recombinant fusion protein of human SLC27A2 (NP_001153101.1).

Conjugation: Unconjugated

Swissprot: O14975

Calculated Molecular Weight: 64 kDa/70 kDa

Observed Molecular Weight: 70 kDa

Concentration: 1 mg/mL

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3

Purification Method: Affinity purification

Application: WB,IHC

Dilution: WB 1:500-1:2000 IHC 1:50-1:200

Storage: Store at -20°C. Avoid freeze / thaw cycles.