003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

PSME3 Polyclonal Antibody

Cat #: G-AB-07897

Host: Rabbit **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:**

Overview:
The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. The immunoproteasome contains an alternate regulator, referred to as the 11S regulator or PA28, that replaces the 19S regulator. Three subunits (alpha, beta and gamma) of the 11S regulator have been identified. This gene encodes the gamma subunit of the 11S regulator. Six gamma subunits combine to form a homohexameric ring. Alternate splicing results in multiple transcript

Gene ID: 10197

Accession #:

Immunogen: A synthetic peptide of human PSME3 (NP_005780.2).

Conjugation: Unconjugated

Swissprot: P61289

Calculated Molecular Weight:

Observed Molecular Weight:

Concentration: 1 mg/mL

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3

Purification Method: Affinity purification

Application: IF

Dilution: IF 1:50-1:200

Storage: Store at -20°C. Avoid freeze / thaw cycles.