003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

PSMD1 Polyclonal Antibody

Cat #: G-AB-09422

Host: Rabbit **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:**

Overview:
The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes the largest non-ATPase subunit of the 19S regulator lid, which is responsible for substrate recognition and binding. Alternatively spliced transcript variants have been found for this gene.

Gene ID: 5707

Accession #:

Immunogen: Recombinant fusion protein of human PSMD1 (NP_002798.2).

Conjugation: Unconjugated

Swissprot: Q99460

Calculated Molecular Weight: 102 kDa/105 kDa

Observed Molecular Weight: 106 kDa

Concentration: 1 mg/mL

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3

Purification Method: Affinity purification

Application: WB

Dilution: WB 1:500-1:2000

Storage: Store at -20°C. Avoid freeze / thaw cycles.