003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

Cyclin D2 Polyclonal Antibody

Cat #: G-AB-09718

Host: Rabbit **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:**

Overview:
The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK4 or CDK6 and functions as a regulatory subunit of the complex, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. Knockout studies of the homologous gene in mouse suggest the essential roles of this gene in ovarian granulosa and germ cell proliferation. High level expression of this gene was observed in ovarian and testicular tumors. Mutations in this gene are associated with megalencephaly-polymicrogyria-polydactyly-hydrocephalus syndrome 3 (MPPH3).

Gene ID: 894

Accession #:

Immunogen: Recombinant fusion protein of human Cyclin D2 (NP_001750.1).

Conjugation: Unconjugated

Swissprot: P30279

Calculated Molecular Weight:

Observed Molecular Weight:

Concentration: 1 mg/mL

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3

Purification Method: Affinity purification

Application: IHC,IF

Dilution: IHC 1:50-1:100 IF 1:50-1:100

Storage: Store at -20°C. Avoid freeze / thaw cycles.