003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

Acetyl-Histone H4-K16 Polyclonal Antibody

Cat #: G-AB-10575

Host: Rabbit **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:**

Overview:
Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in a histone cluster on chromosome 1. This gene is one of four histone genes in the cluster that are duplicated; this record represents the centromeric copy.

Gene ID: 8370

Accession #:

Immunogen: A synthetic acetylated peptide around K16 of human Histone H4 (NP_001029249.1).

Conjugation: Unconjugated

Swissprot: P62805

Calculated Molecular Weight: 11 kDa

Observed Molecular Weight: 13 kDa

Concentration: 1 mg/mL

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3

Purification Method: Affinity purification

Application: WB,IF

Dilution: WB 1:500-1:2000 IF 1:50-1:200

Storage: Store at -20°C. Avoid freeze / thaw cycles.