003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

# RPLP0 Polyclonal Antibody

**Cat #:** G-AB-10714

**Host:** Rabbit **Reactivity:** Human, Mouse

**Isotype:** IgG **Clonality:**

**Overview:**
Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein, which is the functional equivalent of the E. coli L10 ribosomal protein, belongs to the L10P family of ribosomal proteins. It is a neutral phosphoprotein with a C-terminal end that is nearly identical to the C-terminal ends of the acidic ribosomal phosphoproteins P1 and P2. The P0 protein can interact with P1 and P2 to form a pentameric complex consisting of P1 and P2 dimers, and a P0 monomer. The protein is located in the cytoplasm. Transcript variants derived from alternative splicing exist; they encode the same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

**Gene ID:** 6175

**Accession #:**

**Immunogen:** Recombinant fusion protein of human RPLP0 (NP_000993.1).

**Conjugation:** Unconjugated

**Swissprot:** P05388

**Calculated Molecular Weight:**

**Observed Molecular Weight:**

**Concentration:** 1 mg/mL

**Buffer:** PBS with 0.02% sodium azide, 50% glycerol, pH7.3

**Purification Method:** Affinity purification

**Application:** IF

**Dilution:** IF 1:50-1:200

**Storage:** Store at -20°C. Avoid freeze / thaw cycles.