003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

PSMB8 Polyclonal Antibody

Cat #: G-AB-11640

Host: Rabbit **Reactivity:** Human, Mouse, Rat

Isotype: IgG **Clonality:**

Overview:
The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. This gene is located in the class II region of the MHC (major histocompatibility complex). Expression of this gene is induced by gamma interferon and this gene product replaces catalytic subunit 3 (proteasome beta 5 subunit) in the immunoproteasome. Proteolytic processing is required to generate a mature subunit. Two alternative transcripts encoding two isoforms have been identified; both isoforms are processed to yield the same mature subunit.

Gene ID: 5696

Accession #:

Immunogen: Recombinant fusion protein of human PSMB8 (NP_004150.1).

Conjugation: Unconjugated

Swissprot: P28062

Calculated Molecular Weight:

Observed Molecular Weight:

Concentration: 1 mg/mL

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3

Purification Method: Affinity purification

Application: IHC,IF

Dilution: IHC 1:50-1:200 IF 1:50-1:200

Storage: Store at -20°C. Avoid freeze / thaw cycles.