003216589045 (EU)
0203393 8531 (UK)
(408)780-0908 (US)
info@gentaur.com
www.caslab.com

EXOSC4 Polyclonal Antibody

Cat #: G-AB-12338

Host: Rabbit **Reactivity:** Human, Rat

Isotype: IgG **Clonality:**

Overview:
Non-catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease activity and participates in a multitude of cellular RNA processing and degradation events. In the nucleus, the RNA exosome complex is involved in proper maturation of stable RNA species such as rRNA, snRNA and snoRNA, in the elimination of RNA processing by-products and non-coding 'pervasive' transcripts, such as antisense RNA species and promoter-upstream transcripts (PROMPTs), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig class switch recombination (CSR) and/or Ig variable region somatic hypermutation (SHM) by targeting AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements (AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. The catalytic

Gene ID: 54512

Accession #:

Immunogen: Recombinant fusion protein of human EXOSC4 (NP_061910.1).

Conjugation: Unconjugated

Swissprot: Q9NPD3

Calculated Molecular Weight: 26 kDa

Observed Molecular Weight: 26 kDa

Concentration: 1 mg/mL

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3

Purification Method: Affinity purification

Application: WB

Dilution: WB 1:500-1:2000

Storage: Store at -20°C. Avoid freeze / thaw cycles.